

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The sums of the horizontal and vertical projections of AB, BC, CD give

8
$$\cos \alpha + 10 \cos \beta + 12 \cos \gamma = 20$$
,
8 $\sin \alpha + 10 \sin \beta - 12 \sin \gamma = 4$.

These six equations furnish the theoretical solution of the problem.

The Proposer furnished a complete solution.

348 (Mechanics). Proposed by ALTON L. MILLER, Ann Arbor, Michigan.

If equilateral triangles be constructed on the sides of any triangle, their centers are the vertices of a new equilateral triangle. Show that the center of gravity of this new equilateral triangle coincides with the center of gravity of the original triangle.

SOLUTION BY EMMA M. GIBSON, Springfield, Mo.

Let ABC be the given triangle and let the coördinates of the vertices A, B, C referred to the rectangular axes ox and oy be (c, o), (o, b), (a, o), respectively. The equation of the line through (a, o) and (o, b) is

 $y = -\frac{b}{a}x + b. ag{1}$

The line from D, the third vertex of the equilateral triangle on BC, through (a/2, b/2) and perpendicular to (1) is

 $y = \frac{a}{b}x + \frac{b^2 - a^2}{2b} \tag{2}$

The line through C making an angle of 60° with (1) is

$$y = \frac{b + a\sqrt{3}}{b\sqrt{3} - a}(x - a). \tag{3}$$

Solving equations (2) and (3), the values of the coördinates of D are found to be $[(a + b\sqrt{3})/2, (b + a\sqrt{3})/2]$.

Similarly the coördinates of F and E are found to be $[(a+c)/2, \sqrt{3}(c-a)/2]$ and $[(c-b\sqrt{3})/2, (b-c\sqrt{3})/2]$, respectively.

Now the centers H, G, I of the three equilateral triangles are

$$\left(\frac{3a+b\sqrt{3}}{6}, \frac{3b+a\sqrt{3}}{6}\right), \quad \left(\frac{a+c}{2}, \frac{\sqrt{3}(c-a)}{6}\right), \quad \left(\frac{3c-b\sqrt{3}}{6}, \frac{3b-c\sqrt{3}}{6}\right),$$

respectively, since $\bar{x} = \frac{1}{3}(x_1 + x_2 + x_3)$, $\bar{y} = \frac{1}{3}(y_1 + y_2 + y_3)$. These points are the vertices of the new triangle and by the formula for the length of a line between two points, the three sides are proved equal. Hence, the new triangle is equilateral.

The coördinates of the center of gravity of the original triangle are $\bar{x} = \frac{1}{3}[a+c]$, $\bar{y} = \frac{1}{3}b$. The coördinates of the center of gravity of the new triangle are

$$\bar{x} = \frac{1}{3} \left[\frac{a+b\sqrt{3}}{2} + \frac{a+c}{2} + \frac{c-b\sqrt{3}}{2} \right] = \frac{1}{3}(a+c),$$

$$\bar{y} = \frac{1}{3} \left[\frac{b+a\sqrt{3}}{2} + \frac{\sqrt{3}(c-a)}{2} + \frac{b-c\sqrt{3}}{2} \right] = \frac{1}{3}b,$$

which are the same as those obtained for the original triangle.

Also solved by Horace Olson and Roger Johnson.

268 (Number Theory). Proposed by FRANK IRWIN, University of California.

Show that in any arithmetical progression, whose first term a_1 and common difference d are positive integers, any required number of consecutive terms may be found, no one of which is a prime number.